



Review

Retention models for ionizable compounds in reversed-phase liquid chromatography

Effect of variation of mobile phase composition and temperature

Martí Rosés*, Xavier Subirats, Elisabeth Bosch

Departament de Química Analítica, Universitat de Barcelona, Martí i Franquès 1-11, E-08028 Barcelona, Spain

ARTICLE INFO

Article history:

Available online 25 December 2008

Keywords:

Reviews
Retention models
Ionizable compounds
Solvation parameter model
Polarity parameter model
Mobile phase composition
Mobile phase pH
Mobile phase temperature

ABSTRACT

General models in reversed-phase liquid chromatography that have been extended to relate retention of ionizable compounds to mobile phase composition, pH and/or temperature are reviewed. In particular, the fundamentals and applications of the solvation parameter model, the polarity parameter model and several classical models based on empirical equations are presented and compared. A main parameter in all these models is the degree of ionization of the acid–base compound, which depends on both the pH of the mobile phase and the acid–base constant of the compound. Thus, on one hand, the different procedures for pH measurement in the mobile phase and their influence on the performance of the models are outlined. On the other hand, equations that relate the variation of the pH of the buffer and the pK_a of the compound with the mobile phase composition and/or temperature are reviewed and their applicability to the retention models critically discussed.

© 2009 Elsevier B.V. All rights reserved.

Contents

1. Introduction	1756
2. Retention models for ionizable solutes	1757
2.1. Empirical models	1758
2.2. Solvation parameter models	1759
2.3. Polarity parameter models	1761
3. Ionization of acid–base compounds in high-performance liquid chromatography mobile phases	1762
3.1. pH measurement in reversed-phase liquid chromatography	1763
3.2. pK_a variation of acid/base analytes with the addition of acetonitrile or methanol	1764
3.3. pH variation of the mobile phase with the addition of acetonitrile or methanol	1765
3.4. Effect of temperature on ionization and retention	1765
List of symbols	1773
References	1774

1. Introduction

The chromatographic retention of ionizable compounds is strongly dependent on the pH of the mobile phase. In a reversed-phase liquid chromatographic system (RPLC), neutral compounds are much more retained than ionic compounds. For an ionizable analyte, i.e. a compound with acid–base properties in the working range of pH, the ratio between the concentrations of neutral

and ionic forms, and therefore chromatographic retention, depends on the pH of the mobile phase and on the pK_a of the compound. This rate can be easily calculated from the well-known mathematical definitions of the acid–base equilibrium constants, which for a monoprotic acid (HA) or base (A) takes the form:



where the electrical charges of HA and A species have been omitted for simplicity. If the compound is a neutral acid, HA has no charge, whereas A has a negative charge (A^-), so the latter it is usually poorly retained as compared with the former. But if the studied

* Corresponding author. Tel.: +34 93 403 92 75; fax: +34 93 402 12 33.
E-mail address: marti.roses@ub.edu (M. Rosés).